



Tipični simptomi / Typical symptoms

- zvižduci / wheezing
- kratkoća daha (zaduha) / shortness of breath (dyspnoea)
- pritisak u prsištu / chest pain
- kašalj / cough
- tipično noću i pred jutro / at night and early morning
- u naporu / during exertion
- potaknuti alergenima i/ili triggered by allergens and/or
- iritansima /irritans

Atipični simptomi / atypical symptoms

- izolirani suhi kašalj / isolated dry cough

RTG srca i pluća (30 p/y -> LDCT) / Chest X-ray (30 p/y -> LDCT)

patološki nalaz / pathological finding
 obrada /work-up

Opstruktivne smetnje ventilacije (FEV1 <80%, FEV1/FVC <0,75-0,80) / Obstructive airflow pattern (FEV1 <80%, FEV1/FVC <0,75-0,80)
 Pozitivan bronhodilatacijski test sa salbutamolom (FEV1: +12% i +200 mL) / Positive bronchodilator test with salbutamol (FEV1: +12% i +200 mL)

DA / YES
 liječiti astmu / treat asthma

Praćenje varijabilnosti PEF-a tijekom 2 tjedna / PEF variability monitoring during 2 weeks
 ili / or
 Pozitivan bronhoprovokacijski test (↓ FEV1 ≥20%) / Positive bronchoprovocation test (↓ FEV1 ≥20%)

DA / YES
 liječiti astmu / treat asthma

eozinofili u iskašljaju i perifernoj krvi / eosinophils present in sputum and blood
 kožni "prick" test ili sIgE i IgE / dermal "prick" test or sIgE and IgE
 FeNO
 pregled ORL-oga / ENT specialist exam

isključiti druge uzroke suhog kašlja / exclude other dry cough etiologies

uputiti na specijalističku obradu / refer to specialist work-up

uvesti niske doze ICS-a / start low dose ICS

liječiti astmu / treat asthma

nema terapijskog odgovora / no response to therapy